

Pelvic Floor Dysfunction

For most people, having a bowel movement is a seemingly automatic function. For some individuals, the process of evacuating stool may be difficult. Symptoms of pelvic floor dysfunction include constipation and the sensation of incomplete emptying of the rectum when having a bowel movement. Incomplete emptying may result in the individual feeling the need to attempt a bowel movement several times within a short period of time. Residual stool left in the rectum may slowly seep out of the rectum leading to reports of bowel incontinence.

The process of defecation (having a bowel movement) requires the coordinated effort of different muscles. The pelvic floor is made up of several muscles that support the rectum like a hammock. When an individual wants to have a bowel movement the pelvic floor muscles are supposed to relax, allowing the rectum to empty. While the pelvic floor muscles are relaxing, muscles of the abdomen contract to help push the stool out of the rectum. Individuals with pelvic floor dysfunction have a tendency to contract instead of relax the pelvic floor muscles. When this happens during an attempted bowel movement, these individuals are effectively pushing against an unyielding muscular wall.

The diagnosis of pelvic floor disorder starts with a careful history regarding an individual's symptoms, medical problems and a history of physical or emotional trauma that may be contributing to their problem. Next the physician examines the patient to identify any physical abnormality. A defecating proctogram is a study commonly used to demonstrate the functional problem in a person with pelvic floor dysfunction. During this study, the patient is given an enema of a thick liquid that can be detected with x-ray. A special x-ray video records the movement of the pelvic floor muscles and the rectum while the individual attempts to empty the liquid from the rectum. Normally the pelvic floor relaxes allowing the rectum to straighten and the liquid to pass out of the rectum. This study will demonstrate if the pelvic floor muscles are not relaxing appropriately and preventing passage of the liquid.

The defecating proctogram is also useful to show if the rectum is folding in on itself (rectal prolapse). Many women have outpouching of the rectum known as a rectocele. Usually a rectocele does not affect the passage of stool. In some instances, however, stool may become trapped in a rectocele causing symptoms of incomplete evacuation. The defecating proctogram helps to identify if liquid is getting trapped in a rectocele when the individual is trying to empty the rectum.

Pelvic floor dysfunction due to non-relaxation of the pelvic floor muscles may be treated with specialized physical therapy known as biofeedback. With biofeedback, a therapist helps to improve a person's rectal sensation and pelvic floor muscle coordination. There are various effective techniques used in biofeedback. Some therapists train patients by teaching them to expel a small balloon placed in the rectum. Another technique uses a small probe placed in the rectum or vagina or electrodes placed on the surface of the skin around the opening to the rectum (anus) and on the abdominal wall. These instruments detect when a muscle is contracting or relaxing and provide visual feedback of the muscle

action. This visual feedback helps the individual to understand the muscle movement and aids in improving muscle coordination. Approximately 75% of individuals with pelvic floor dysfunction experience significant improvement with biofeedback.

Abnormalities identified with a defecating proctogram such as rectal prolapse and rectocele may be treated with a surgical procedure.

What is a Colorectal Surgeon?

Colon and rectal surgeons are experts in the surgical and non-surgical treatment of colon and rectal problems. They have completed advanced training in the treatment of colon and rectal problems in addition to full training in general surgery. Colon and rectal surgeons treat benign and malignant conditions, perform routine screening examinations and surgically treat problems when necessary.