COLONOSCOPY PREP INSTRUCTIONS (AM exam) Miralax (same as Glycolax)

You will need to purchase these items below-these are all over the counter and you can find listed options of alternatives on the back of this sheet.

Miralax bottle (238 grams or 8.3 ounces)

(4) Dulcolax or bisacodyl tablets (5 milligram tablets)

64 oz. of re-hydration (sports) drink except no red drinks (See back of this sheet for list of brands to choose from)

<u>Diff OND</u> - one day prior to colonoscopy appointment	
7:00 am	Begin clear liquid diet (list on back). Absolutely NO solid
	food or alcohol
10:00 am	Take 2 bisacodyl tablets with water
3:00 pm	Take remaining 2 bisacodyl tablets with water (do not
	crush or chew and do not take within 1 hour of taking
	an antacid
4:00 pm	Combine the 238 gram bottle (or 8.3oz.) of Miralax
	powder into an empty pitcher along with 64 oz. of
	rehydration (sports) drink. Mix and chill if desired.
5:00 pm	Start to drink the Miralax. Drink (1) 8oz. glass every 15
	minutes over a two hour period. Drink each glass
	quickly rather than drinking small amounts
	continuously. Continue to drink clear liquids the
	remainder of the evening.

<u>DAY ONE</u> = one day prior to colonoscopy appointment

<u>DAY TWO</u> = day of colonoscopy exam

1.	Continue to drink clear liquids until 2 hours before your colonoscopy. 2 hours before your colonoscopy, stop ALL fluids and take nothing (NO FOOD IR DRINK) by mouth.
2.	For use of your medications, refer to orange medication instruction sheet enclosed.
3.	Call (585-461-2498) if you have any questions regarding any of these instructions.
4.	Arrive at your assigned time.

****We encourage clear liquid intake up to 2 hours before your procedure time****

NO SEEDS, NUTS, POPCORN, CORN OR QUINOA 3 DAYS PRIOR TO YOUR PROCEDURE

<u>CLEAR LIQUIDS (NO ALCOHOL)</u>: *Do not drink any liquid that is **RED***

- Water
- Clear juices (apple, white grape juice, orange drink, white cranberry juice)
- Black coffee or tea (no creamers) sugar OK
- Bouillon or clear broth
- Soda pop (Sprite, Ginger-ale, Mountain Dew), Gatorade (no red)
- Jell-o (no red), popsicles (no red), Lemon Ice

BOWEL PREP TABLETS: (any of these brands)

These are the PILLS to take at the start of your bowel prep

- Bisacodyl (Generic)
- Dulcolax
- Dulcogen
- Alophen

DO NOT USE STOOL SOFTENERS

- Bisco-lax
- Carters Little Pills
- Colax
- Fleet Bisacodyl
- Modane

RE-HYDRATION (SPORT) DRINKS: 64 oz. *NO RED DRINKS*

This is the liquid used to mix with the powder (below)

- Powerade (contains less sugar than Gatorade)
- Powerade zero (even less sugar, best for diabetics)
- G2 (same amount of electrolytes as Gatorade, just less calories)
- MVP sports drink (Wegmans brand)
- MVP 2 sports drink (Wegmans brand) best for diabetics
- AllSport
- HydraLife powder
- Gatorade

BOWEL PREP POWDER

This is the powder to mix with the liquid (above)

- Glycolax
- PureLax
- ClearLax
- Miralax

Please Note:

- Please wear socks as it sometimes can get cool in the procedure room.
- No nail polish on fingers only (you do not need to remove acrylic nails). The nail polish may interfere with the oxygen sensor that is placed on your finger.
- Diabetics may use Powerade Zero as it contains less sugar than regular Powerade.
- Please leave all valuables at home. Do not wear jewelry/watches to procedure.